

On the solutions to accelerating cosmologies

Masato Ito *

Department of Physics, Nagoya University, Nagoya, JAPAN 464-8602

Abstract

Motivated by recent accelerating cosmological model, we derive the solutions to vacuum Einstein equation in $(d + 1)$ -dimensional Minkowski space with n -dimensional hyperbolic manifold. The conditions of accelerating expansion are given in such a set up.

*E-mail address: mito@eken.phys.nagoya-u.ac.jp

Recently, Townsend and Wohlfarth discovered a solution of the four-dimensional accelerating cosmologies with compact hyperbolic manifold [1]. By avoiding the assumptions of a no-go theorem [2], it was pointed out that the accelerating phase arises in a interval. The features of the model are that the four-dimensional metric is Einstein frame and the compact hyperbolic manifold is time-dependent. Before [1], the investigations of FRW universes with closed hyperbolic internal space have been widely performed [3, 4, 5, 6, 7]. Furthermore it has been shown that the model is closely connection with the S-branes which correspond to time-dependent solutions of supergravity with p -form [8, 9, 10, 11, 12, 13, 14]. Thus it is expected that the accelerating universe supported by astronomical observations may be explained by string/M-theory.

From the theoretical point of view, we are interested in the higher dimensional cosmological model motivated by the model of Ref.[1]. In this letter, we derive the solutions to the vacuum Einstein equation in time-dependent $\mathcal{M}_{d+1} \times H_n$, where $\mathcal{M}_{d+1}$ and H_n are $(d+1)$ -dimensional Minkowski and n -dimensional hyperbolic manifold, respectively. Furthermore we investigate the conditions of accelerated expansion of the present model.

Adopting the Einstein frame metric for $d \geq 3$, we can take ansatz for time-dependent metric

$$ds^2 = b^{\frac{4}{d-1}}(t)ds_{d+1}^2 + b^{-\frac{4}{n}}(t)d\hat{s}_n^2, \quad (1)$$

where

$$ds_{d+1}^2 = -a^{2d}(t)dt^2 + a^2(t)dx_d^2, \quad (2)$$

where $a(t)$ corresponds to the scale factor of d -dimensions. For the n -dimensional hyperbolic manifold, the Ricci tensor is given by

$$\hat{\mathcal{R}}_{ab} = -(n-1)g_{ab}. \quad (3)$$

The Ricci tensors for metric (1) are evaluated as follows,

$$\begin{aligned} \mathcal{R}_{tt} &= -\frac{2}{d-1}\frac{\ddot{b}}{b} - 2\left(\frac{1}{d-1} + \frac{2}{n}\right)\frac{\dot{b}^2}{b^2} - d\frac{\ddot{a}}{a} + d^2\frac{\dot{a}^2}{a^2} \\ \mathcal{R}_{\mu\nu} &= a^{2(1-d)}\left\{\frac{2}{d-1}\left(\frac{\dot{b}}{b}\right) + \left(\frac{\dot{a}}{a}\right)\right\}g_{\mu\nu} \end{aligned}$$

$$\mathcal{R}_{ab} = \left[-\frac{2}{n} b^{-4(\frac{1}{n} + \frac{1}{d-1})} a^{-2d} \left(\frac{\dot{b}}{b} \right) - (n-1) \right] g_{ab} \quad (4)$$

where μ, ν and a, b represent $\mathcal{M}_{d+1}$ and H_n , respectively. The dot denotes the derivative with respect to the time t . In order to simplify the vacuum Einstein equation for metric ansatz, we define variables X and Y by

$$\begin{aligned} b &= e^X \\ a &= e^Y. \end{aligned} \quad (5)$$

Using the new variables, the equations of motion can be rewritten as

$$-\frac{2}{d-1} \ddot{X} - 4 \left(\frac{1}{d-1} + \frac{2}{n} \right) \dot{X}^2 - d \ddot{Y} + d(d-1) \dot{Y}^2 = 0, \quad (6)$$

$$\frac{2}{d-1} \ddot{X} + \ddot{Y} = 0, \quad (7)$$

$$\ddot{X} + \frac{1}{2} n(n-1) e^{4(\frac{1}{d-1} + \frac{1}{n})X + 2dY} = 0. \quad (8)$$

Performing the integral of (7), and we have

$$\frac{2}{d-1} X + Y = ft + g, \quad (9)$$

where f and g are integration constants. Substituting (9) into (6) and (8), the equations of motion can be written in terms of X

$$\ddot{X} + 2 \frac{n-1}{n} \dot{X}^2 - 2fd\dot{X} + \frac{1}{2} d(d-1)f^2 = 0 \quad (10)$$

$$\ddot{X} = -\frac{1}{2} n(n-1) e^{2gd} e^{-4\frac{n-1}{n}X + 2dft}, \quad (11)$$

In order to solve (11), we define Z by

$$Z = -4 \frac{n-1}{n} X + 2dft. \quad (12)$$

Consequently, the second order differential equation can be obtained

$$\ddot{Z} = 2(n-1)^2 e^{2gd} e^Z. \quad (13)$$

Performing the integral of the above equation, we have

$$\dot{Z}^2 = 4(n-1)^2 e^{2gd} e^Z + h, \quad (14)$$

where h is positive integration constant. Therefore, the solution to the Z is given by

$$e^Z = \frac{h}{4(n-1)^2} \frac{e^{-2gd}}{\sinh^2 \frac{\sqrt{h}}{2} (t+t_0)}, \quad (15)$$

where t_0 is the integration constant. Furthermore, (12) can be rewritten in terms of Z

$$2n\ddot{Z} - n\dot{Z}^2 + 4f^2d(n+d-1) = 0. \quad (16)$$

Substituting (13) and (14) into (16), the relation between the integration constants h and f can be expressed as

$$h = 4 \frac{d(n+d-1)}{n} f^2. \quad (17)$$

Here we simply take $f = 1$, $g = 0$, $t_0 = 0$. Combining with (9), (12) and (15), and we have

$$b = e^X = e^{\frac{nd}{2(n-1)}t} \left(\frac{\gamma}{(n-1) \sinh \gamma|t|} \right)^{-\frac{n}{2(n-1)}} \quad (18)$$

$$a = e^Y = e^{-\frac{n+d-1}{(n-1)(d-1)}t} \left(\frac{\gamma}{(n-1) \sinh \gamma|t|} \right)^{\frac{n}{(d-1)(n-1)}} \quad (19)$$

where

$$\gamma = \sqrt{\frac{d(n+d-1)}{n}}. \quad (20)$$

Note that the argument of the hyperbolic sine can be taken to be positive in order that the solutions are real.

For $d = 3$, obviously, (18) and (19) are completely consistent with the metric in Ref.[1]. Furthermore it is considered that the results obtained here are included in S-brane solutions in arbitrary dimensions derived by N. Ohta [10].

Below we shall investigate expansion and acceleration in the case of $d \geq 3$. From the metric (2), the proper time η for $(d+1)$ -dimensional world is given by $d\eta = a^d dt$. The condition of expanding $(d+1)$ -dimensional universe, $da/d\eta > 0$, leads to

$$M(t) = 1 + \sqrt{\frac{nd}{n+d-1}} \coth \left(\sqrt{\frac{d(n+d-1)}{n}} t \right) < 0. \quad (21)$$

The above inequality is satisfied for negative t since the coefficient of hyperbolic cotangent is always larger than one.

Moreover the condition of accelerating $(d + 1)$ -dimensional universe, $d^2 a/d\eta^2 > 0$, leads to

$$M^2(t) < \frac{d(n-1)}{n+d-1} \frac{1}{\sinh^2 \left(\sqrt{\frac{d(n+d-1)}{n}} t \right)}. \quad (22)$$

The accelerating phase arises if the conditions (21) and (22) are simultaneously satisfied at negative t . As for the asymptotic relationship between time t and proper time η , note that $t \rightarrow -\infty$ corresponds to $\eta \rightarrow +0$ while $t \rightarrow -0$ corresponds to $\tau \rightarrow \infty$. Furthermore the asymptotic behavior of scale factor for η is expressed as

$$a(\eta) \sim \begin{cases} \eta^{\frac{1}{d}} & \text{for } \eta \rightarrow +0 \ (t \rightarrow -\infty) \\ \eta^{\frac{n}{n+d-1}} & \text{for } \eta \rightarrow \infty \ (t \rightarrow -0) \end{cases}. \quad (23)$$

For small η , note that scale factor is independent of n .

The interval of accelerating expansion is determined by $d^2 a/d\eta^2 = 0$. Therefore we get

$$\begin{aligned} t_1 &= -\sqrt{\frac{n}{d(n+d-1)}} \log \frac{\sqrt{n-1}(\sqrt{d}+1)}{\sqrt{nd}-\sqrt{n+d-1}} \\ t_2 &= -\sqrt{\frac{n}{d(n+d-1)}} \log \frac{\sqrt{n-1}(\sqrt{d}-1)}{\sqrt{nd}-\sqrt{n+d-1}}, \end{aligned} \quad (24)$$

where t_1 and t_2 are start and end of acceleration, respectively. Using the above results, we can estimate the e-foldings number in an interval of acceleration, namely, $\log(a(t_2)/a(t_1)) = \mathcal{O}(1)$. As indicated in Ref.[15], since the e-foldings number is of order one, it is questionable whether the present model is worthy of inflation model. It is expected that models or mechanisms for enlarging e-foldings number are proposed in the future.

In summary, we derived the solutions to vacuum Einstein equation in $\mathcal{M}_{d+1} \times H_n$ with Einstein frame metric for $d \geq 3$, triggered by recent accelerating cosmological model of Ref.[1]. We provided the conditions of accelerating expansion and the asymptotic behaviors of scale factor. Moreover, the start and end of acceleration are given and it was pointed out that the e-foldings number during the interval is approximately of order one for proper values of n, d .

Acknowledgements:

We thank N. Ohta for pointing out errors in the earlier version of the paper and useful correspondence.

Note added: After we submitted this paper to the hep archives, we learned the higher dimensional accelerating cosmologies with product spaces proposed by C-M. Chen et al.[15]. It was pointed out that the model considered here is related to the S-branes by N. Ohta [10].

References

- [1] P. K. Townsend and M. N. Wohlfarth, “Accelerating cosmologies from compactification,” arXiv:hep-th/0303097.
- [2] J. M. Maldacena and C. Nunez, “Supergravity description of field theories on curved manifolds and a no go theorem,” Int. J. Mod. Phys. A **16**, 822 (2001) [arXiv:hep-th/0007018].
- [3] G. D. Starkman, D. Stojkovic and M. Trodden, “Large extra dimensions and cosmological problems,” Phys. Rev. D **63**, 103511 (2001) [arXiv:hep-th/0012226].
- [4] N. Kaloper, J. March-Russell, G. D. Starkman and M. Trodden, “Compact hyperbolic extra dimensions: Branes, Kaluza-Klein modes and cosmology,” Phys. Rev. Lett. **85**, 928 (2000) [arXiv:hep-ph/0002001].
- [5] G. D. Starkman, D. Stojkovic and M. Trodden, “Homogeneity, flatness and ‘large’ extra dimensions,” Phys. Rev. Lett. **87**, 231303 (2001) [arXiv:hep-th/0106143].
- [6] P. J. Silva, “Stabilization of hyperbolic brane-world scenarios,” arXiv:hep-th/0202160.
- [7] S. Nasri, P. J. Silva, G. D. Starkman and M. Trodden, “Radion stabilization in compact hyperbolic extra dimensions,” Phys. Rev. D **66**, 045029 (2002) [arXiv:hep-th/0201063].
- [8] M. Gutperle and A. Strominger, “Spacelike branes,” JHEP **0204**, 018 (2002) [arXiv:hep-th/0202210].

- [9] M. Kruczenski, R. C. Myers and A. W. Peet, “Supergravity S-branes,” JHEP **0205**, 039 (2002) [arXiv:hep-th/0204144].
- [10] N. Ohta, “Intersection rules for S-branes,” Phys. Lett. B **558**, 213 (2003) [arXiv:hep-th/0301095].
- [11] N. Ohta, “Accelerating cosmologies from S-branes,” arXiv:hep-th/0303238.
- [12] N. Ohta, “A study of accelerating cosmologies from superstring / M theories,” arXiv:hep-th/0304172.
- [13] R. Emparan and J. Garriga, “A note on accelerating cosmologies from compactifications and S-branes,” arXiv:hep-th/0304124.
- [14] M. N. Wohlfarth, “Accelerating cosmologies and a phase transition in M-theory,” arXiv:hep-th/0304089.
- [15] C. M. Chen, P. M. Ho, I. P. Neupane and J. E. Wang, “A note on acceleration from product space compactification,” arXiv:hep-th/0304177.